

FIRST COUNTABLE SPACES WITHOUT POINT-COUNTABLE π -BASES

ISTVÁN JUHÁSZ, LAJOS SOUKUP, AND ZOLTÁN SZENTMIKLÓSSY

ABSTRACT. We answer several questions of V. Tkačuk from [*Point-countable π -bases in first countable and similar spaces*, Fund. Math. 186 (2005), pp. 55–69.] by showing that

- (1) there is a ZFC example of a first countable, 0-dimensional Hausdorff space with no point-countable π -base (in fact, the order of any π -base of the space is at least $\aleph_\omega$);
- (2) if there is a κ -Suslin line then there is a first countable GO space of cardinality κ^+ in which the order of any π -base is at least κ ;
- (3) it is consistent to have a first countable, hereditarily Lindelöf regular space having uncountable π -weight and ω_1 as a caliber (of course, such a space cannot have a point-countable π -base).

1. INTRODUCTION

V. Tkačuk in [8] has recently proved under CH that any first countable Hausdorff space that is Lindelöf or CCC has a point-countable π -base. (Actually, in [8] all spaces are assumed to be Tychonov, but the proof only needs Hausdorff.) Tkačuk's motivation was to extend (at least partially) Šapirovskii's celebrated ZFC result saying that any countably tight compactum has a point-countable π -base, from compact spaces to Lindelöf ones. So it was natural to ask if his use of CH was necessary. To Tkačuk's amazement (and embarrassment), he could come up with no example of a first countable space without a point-countable π -base, not even consistently.

Our aim here is to remedy this embarrassing situation and provide not only ZFC (and several consistent) examples of first countable (Tychonov) spaces without point-countable π -bases but also examples which show that, indeed, Tkačuk's CH results cannot be proved in ZFC

2000 *Mathematics Subject Classification.* 54A25, 54A35, 54D20, 54D70, 54F05.

Key words and phrases. First countable space, point-countable π -base, Lindelöf space, CCC space, caliber.

Research supported by OTKA grant no. 61600.

alone. In this manner we succeeded in answering 7 of the 12 questions that were listed at the end of [8].

In what follows, we shall use the notation and terminology of [4]. In particular, $\pi sw(X)$ denotes the π -separating weight of X , that is the minimum order of a π -base of the space X , see p. 74 of [4]. Note that $\pi sw(X) \leq \omega$ is then equivalent to the statement: X has a point-countable π -base.

2. ZFC EXAMPLES

The key to Tkačuk's above mentioned CH results in [8] was his theorem 3.1 which says that if X has countable tightness and π -character, moreover $d(X) \leq \omega_1$ then $\pi sw(X) \leq \omega$. In his list of problems (problem 4.11), Tkačuk himself asked if the assumption of countable tightness could be omitted here. It is immediate from our next result that this question has an affirmative answer.

Theorem 1. *Let X be any topological space with $d(X) \leq \pi\chi(X)^+$. Then $\pi sw(X) \leq \pi\chi(X)$.*

Proof. Let us set $\pi\chi(X) = \kappa$. If $d(X) \leq \kappa$ then we even have $\pi(X) = \kappa$. So we may assume $d(X) = \kappa^+$ and, as is well-known, we may then fix a dense set $D = \{x_\alpha : \alpha < \kappa^+\}$ that is left-separated in this well-ordering. This means that for every $\alpha < \kappa^+$ there is a neighbourhood U_α of x_α with

$$\{x_\beta : \beta < \alpha\} \cap U_\alpha = \emptyset.$$

Let us now fix a local π -base $\mathcal{B}_\alpha$ of the point x_α such that $|\mathcal{B}_\alpha| \leq \kappa$ and $B \subset U_\alpha$ whenever $B \in \mathcal{B}_\alpha$. Then $\mathcal{B} = \bigcup \{\mathcal{B}_\alpha : \alpha < \kappa^+\}$ is clearly a π -base of X such that for every $x_\beta \in D$ we have

$$ord(x_\beta, \mathcal{B}) = |\{B \in \mathcal{B} : x_\beta \in B\}| \leq \kappa.$$

We claim that then we have

$$ord(\mathcal{B}) = \sup\{ord(x, \mathcal{B}) : x \in X\} \leq \kappa$$

as well. Assume, on the contrary, that $ord(x, \mathcal{B}) = \kappa^+$ for some point $x \in X$. Since $\pi\chi(x, X) \leq \kappa$, this implies that there are κ^+ many members of $\mathcal{B}$ (containing x) that include a fixed non-empty open set V . This, however, is impossible because $D \cap V \neq \emptyset$. $\square$

We may now turn to our main aim that is to produce, in ZFC, first countable spaces without point-countable π -bases.

Theorem 2. *There is a first countable, 0-dimensional Hausdorff (hence Tychonov) space X with $\pi sw(X) \geq \aleph_\omega$.*

Proof. The underlying set of our space is $X = \prod\{\omega_n : n < \omega\}$. For $f, g \in X$ we write $f \leq g$ to denote that $f(n) \leq g(n)$ for all $n < \omega$. The topology τ that we shall consider on X will be generated by all sets of the form $U_n(f)$ (with $f \in X$ and $n < \omega$), where

$$U_n(f) = \{g \in X : f \leq g \text{ and } f \upharpoonright n = g \upharpoonright n\}.$$

Note that if $g \in U_n(f)$ then $U_n(g) \subset U_n(f)$, and if $g \notin U_n(f)$ then there is a $k < \omega$ such that $U_k(g) \cap U_n(f) = \emptyset$. It follows immediately from this that, for any $f \in X$, the family $\{U_n(f) : n < \omega\}$ forms a clopen neighbourhood base of f with respect to the topology τ , consequently the space $\langle X, \tau \rangle$ is indeed first countable, 0-dimensional, and Hausdorff.

It is also easy to see from the definitions that if $\{U_{n_\alpha}(f_\alpha) : \alpha < \kappa\}$ is a π -base of τ then $\{f_\alpha : \alpha < \kappa\}$ must be cofinal in the partial order $\langle X, \leq \rangle$. But it is well-known that the latter has cofinality greater than $\aleph_\omega$, consequently we have $\pi(X) > \aleph_\omega$. (Actually, it is easy to see that $\pi(X) = \text{cf}(\langle X, \leq \rangle)$ but we shall not need this.)

Next we claim that for any $k < \omega$ the pair $(\aleph_{\omega+1}, \aleph_k)$ is a *pair caliber* of the space X , i. e. among any $\aleph_{\omega+1}$ open sets one can find $\aleph_k$ whose intersection is non-empty. Indeed, we may assume without any loss of generality that we have a family of basic open sets of the form $\{U_n(f) : f \in F\}$ where $F \in [X]^{\aleph_{\omega+1}}$ and $n > k$ is fixed. We may also assume that $f \upharpoonright n = \sigma$ for a fixed $\sigma \in \prod_{i < n} \omega_i$ whenever $f \in F$. Now let $G \subset F$ with $|G| = \aleph_k$ then there is $g \in X$ with $g \upharpoonright n = \sigma$, moreover $f(i) < g(i)$ for all $i \geq n$ and $f \in G$. But then we clearly have $g \in \bigcap \{U_n(f) : f \in G\}$.

Now, putting together the previous two paragraphs we can immediately conclude that the order of any π -base of $\langle X, \tau \rangle$, being of cardinality $> \aleph_\omega$, must be at least $\aleph_\omega$, i. e. $\pi sw(X) \geq \aleph_\omega$. □

The cardinality of our above example is $\aleph_\omega^{\aleph_0}$ that is much larger than the optimal value $\aleph_2$ permitted by theorem 1. So it is natural to raise the question if we could find other examples of smaller cardinality. As it turns out, we can do slightly better by choosing an appropriate subspace Y of the space X from theorem 2. First, however, we need to fix some notation. For $f, g \in X = \prod\{\omega_n : n < \omega\}$ we write $f <^* g$ to denote that $|\{n < \omega : f(n) \geq g(n)\}|$ is finite, i. e. f is below g modulo finite. Similarly, we write $f =^* g$ to denote that $|\{n < \omega : f(n) \neq g(n)\}|$ is finite. Finally, we recall that there is in X a transfinite sequence of order type $\omega_{\omega+1}$ that is increasing with respect to $<^*$.

Theorem 3. *Let $\{f_\alpha : \alpha < \omega_{\omega+1}\} \subset X$ be an increasing sequence with respect to $<^*$ and set*

$$Y = \{f \in X : \exists \alpha < \omega_{\omega+1} \text{ with } f =^* f_\alpha\}.$$

Then the subspace Y of X , with the subspace topology inherited from τ , also satisfies $\pi sw(Y) \geq \aleph_\omega$.

Proof. The proof is very similar to that of theorem 2. First we note that, trivially, again we have $\pi(Y) > \aleph_\omega$. Next, we show that $(\aleph_{\omega+1}, \aleph_k)$ is a pair caliber of Y for each $k < \omega$. To see this, we let again consider any family $\{U_n(f) : f \in F\}$ where $F \in [Y]^{\aleph_{\omega+1}}$ and $n > k > 0$, moreover $f \upharpoonright n = \sigma$ for a fixed $\sigma \in \prod_{i < n} \omega_i$ whenever $f \in F$. Let us choose any subset $G \subset F$ with $|G| = \aleph_k$, then there is an ordinal $\alpha < \omega_{\omega+1}$ such that $g <^* f_\alpha$ for all $g \in G$. Clearly, then we may find an integer $m \geq n$ such that the set

$$G^* = \{g \in G : \forall i \geq m (g(i) < f_\alpha(i))\}$$

also has cardinality $\aleph_k$.

Note that if $n \leq j < m$ then $\{g(j) : g \in G^*\}$ is bounded in ω_j , hence we may find a function $f \in Y$ such that $f \upharpoonright n = \sigma$, if $n \leq j < m$ then $g(j) < f(j)$ for all $g \in G^*$, moreover $f(i) = f_\alpha(i)$ whenever $m \leq i < \omega$. But then we have $f \in \bigcap \{U_n(g) : g \in G^*\} \cap Y$. $\square$

We were unable to produce a ZFC example of a first countable space without a point-countable π -base of cardinality less than $\aleph_{\omega+1}$. This leads us to the following intriguing open question.

Problem 4. *Is there, in ZFC, a first countable (Tychonov) space of cardinality less than $\aleph_\omega$ that has no point-countable π -base?*

Actually, at this point we do not even have such an example of cardinality $\aleph_\omega$. We conjecture, however, that having such an example is equivalent to having one of size $< \aleph_\omega$. In fact, we could verify this conjecture under the assumption $2^{\aleph_1} < \aleph_\omega$.

Theorem 5. *Assume that $2^{\aleph_1} < \aleph_\omega$ and X is a first countable space of cardinality $\aleph_\omega$. If every subspace of X of cardinality $< \aleph_\omega$ has a point-countable π -base then so does X .*

Proof. Let us start by giving a (very natural) definition. A family $\mathcal{B}$ of non-empty open sets in X is said to be an *outer π -base* of a subspace $Y \subset X$ if for every open set U with $U \cap Y \neq \emptyset$ there is a member $B \in \mathcal{B}$ such that $B \subset U$. We claim that, under the assumptions of our theorem, every subspace of X of cardinality $< \aleph_\omega$ actually has a point-countable outer π -base. Thus if we have $X = \bigcup_{n < \omega} Y_n$ where

$|Y_n| < \aleph_\omega$ for all $n < \omega$ and $\mathcal{B}_n$ is a point-countable outer π -base of Y_n in X then $\bigcup_{n < \omega} \mathcal{B}_n$ is clearly a point-countable π -base of X .

To prove the above claim let us consider an ω_1 -closed elementary submodel M with $|M| < \aleph_\omega$ of a "universe" $H(\theta)$. (As usual, here $H(\theta)$ is the collection of all sets of hereditary cardinality $< \theta$, and M is ω_1 -closed means that $[M]^{\leq \omega_1} \subset M$.) The regular cardinal θ is chosen so large that $H(\theta)$ (and also M) contains X and everything else that is relevant, e. g. a map $\mathcal{V}$ that assigns to every point $x \in X$ a countable open neighbourhood base $\mathcal{V}_x$. Clearly, $2^{\aleph_1} < \aleph_\omega$ implies that for every $Y \in [X]^{< \aleph_\omega}$ there is such an elementary submodel M with $Y \subset M$. Consequently, our claim will be proven if we show that $X \cap M$ has a point-countable outer π -base in X whenever M is like above.

To see this, note first that for every point $x \in X \cap M$ we have $\mathcal{V}_x \in M$ and hence $\mathcal{V}_x \subset M$ as well. Consequently $\mathcal{V}_M = \bigcup \{\mathcal{V}_x : x \in X \cap M\} \subset M$ is an outer base of $X \cap M$ in X , hence we may choose a subfamily $\mathcal{B} \subset \mathcal{V}_M$ such that $\mathcal{B} \restriction M = \{B \cap M : B \in \mathcal{B}\}$ is a point-countable π -base of the subspace $X \cap M$.

It suffices to show now that $\mathcal{B}$ is a point-countable outer π -base of $X \cap M$ in X . Indeed, $\mathcal{B}$ is point-countable for if $\mathcal{U} \in [\mathcal{B}]^{\omega_1}$ then we have $\mathcal{U} \in M$ because M is ω_1 -closed, and thus $\bigcap \mathcal{U} \neq \emptyset$ would imply $\bigcap \mathcal{U} \cap M \neq \emptyset$, contradicting that $\mathcal{B} \restriction M$ is point-countable. (Here we also used the fact that, by elementarity, the correspondance $B \mapsto B \cap M$ is one-to-one on $\mathcal{B} \subset M$.) Finally, $\mathcal{B}$ is an outer π -base of $X \cap M$ because if U is open with $x \in U \cap M \neq \emptyset$ then there is $V \in \mathcal{V}_x \subset M$ with $V \subset U$, hence if $B \in \mathcal{B}$ with $B \cap M \subset V \in M$ then we also have $B \subset V \subset U$. $\square$

3. EXAMPLES FROM HIGHER SUSLIN LINES

We start this section by giving a theorem that, not surprisingly, will turn out to be very useful in finding (first countable) spaces without point-countable π -bases.

Theorem 6. *Assume that X is a topological space which has a π -base $\mathcal{B}$ such that $\text{ord}(\mathcal{B})^+ < d(X)$. Then X has a discrete subspace D with $|D| \geq d(X)$.*

Proof. Let us first choose a point $x_B \in B$ from each $B \in \mathcal{B}$. Clearly, then the set $S = \{x_B : B \in \mathcal{B}\}$ is dense in X , hence we have

$$|S| \geq d(X) > \text{ord}(\mathcal{B})^+.$$

We now define a set mapping F on S by the following stipulation: For any point $x \in S$ let us put

$$F(x) = \{x_B \in S : x \in B\} \in [S]^{\leq \text{ord}(\mathcal{B})}.$$

By Hajnal's set mapping theorem (see [2]) then there is a free set $D \subset S$ for the set mapping F with $|D| = |S|$. This means that for every $x \in D$ we have $D \cap F(x) \subset \{x\}$. But every member of D is of the form x_B for some $B \in \mathcal{B}$, and we claim that for this point we have $B \cap D = \{x_B\}$. Indeed, $x_B \in B \cap D$ is obvious, and if $x \in D$ is different from x_B then $x_B \notin F(x)$ implies $x \notin B$. Consequently, D is as required. $\square$

It is an immediate consequence of theorem 6 that a space X satisfying $s(X) < d(X) \geq \omega_2$ cannot have a point-countable π -base. Unfortunately, we do not know if there is in ZFC a first countable Tychonov space like that. (Recall that the solution of Tkačuk's problems from require Tychonov examples.)

If, however, we are satisfied with Hausdorff examples then we are much better off. In fact it was shown in [3] that there is a natural left-separated refinement σ of the euclidean topology τ on the real line $\mathbb{R}$ that is first countable and hereditarily Lindelöf, moreover the density of σ is equal to $\text{cf}(\mathfrak{c})$. Consequently, if $\text{cf}(\mathfrak{c}) > \omega_1$ then, by theorem 6, the space $\langle \mathbb{R}, \sigma \rangle$ has no point-countable π -base. This is of interest because it shows that, at least for Hausdorff spaces, Tkačuk's CH result that was mentioned in the introduction is not valid without CH, for $\langle \mathbb{R}, \sigma \rangle$ is both Lindelöf and CCC. Actually, it is very easy to show that, assuming $\text{cf}(\mathfrak{c}) > \omega_1$ again, something stronger than CCC can be established, namely that ω_1 is a caliber of this space and, for Hausdorff spaces, this settles one more question of Tkačuk from [8]. In the next section we shall produce (consistent) Tychonov examples with these properties but that will require more work.

Next we shall consider higher Suslin lines, these are ordered spaces whose spread is less than their density. More precisely, we shall consider first countable variations of them that retain this property. For different purposes, this construction had been already used in theorem 1.1 of [5], although there CH was additionally assumed.

Let κ be an infinite cardinal. We shall call a continuous linear order $\langle L, < \rangle$, equipped with the order topology generated by $<$, a κ -Suslin line if there are no more than κ disjoint open intervals in L (i. e. $c(L) \leq \kappa$), although the density $d(L)$ of L is bigger than κ . (It is known that the existence of a κ -Suslin line is equivalent to the existence of a κ -Suslin tree, but this will be irrelevant for us.) Thus, an ordinary Suslin line is the same as an ω -Suslin-line and by a higher Suslin line we mean a κ -Suslin line where $\kappa > \omega$.

The main result of this section is the following theorem that, in particular, yields us a consistent example of a first countable GO-space without a point-countable π -base of the minimum possible cardinality

ω_2 . (Recall that GO-spaces, or generalized ordered spaces, are the subspaces of linearly ordered spaces.)

Theorem 7. *If there is a κ -Suslin line $\langle L, < \rangle$ then there is a first countable GO-space X with $|X| = \kappa^+$ and $\pi sw(X) = \kappa$.*

Proof. Let Z be the set of all those points $x \in L$ that have left-character ω , that is the open half line $(\leftarrow, x)$ has cofinality ω with respect to $<$. Since $\langle L, < \rangle$ is continuous, Z is clearly dense in L . It follows that $d(Z) = d(L) = \kappa^+$ because $d(L) \leq c(L)^+$ holds for any linearly ordered space L , see e. g. [1]. Now let X be any dense subspace of Z (and hence of L) with $|X| = \kappa^+$.

We consider X with the left-Sorgenfrey topology σ , i. e. for any $x \in X$ the half-open intervals $(y, x]$ form a σ -local base. Then σ is clearly finer than the order topology on X , hence the density of $\langle X, \sigma \rangle$ must be bigger than κ . It is clear from the definition that σ is a first countable topology.

Also, $\langle X, \sigma \rangle$ is a GO space because it is homeomorphic to the subspace topology on $X \times \{0\}$ inherited from the order topology on $L \times 2$ taken with the lexicographic order. Finally, we clearly have $c(X, \sigma) = c(L)$, moreover $s(X, \sigma) = c(X, \sigma)$ is known to hold for GO-spaces, see e. g. 2.23 of [4]. Consequently we have $s(X) \leq \kappa < d(X)$ and so theorem 6 implies $\pi sw(X) \geq \kappa$. But by theorem 1 we must have then $\pi sw(X) = \kappa$. $\square$

In particular, the existence of an ω_1 -Suslin line implies that of a first countable GO space of cardinality ω_2 without a point-countable π -base.

Finally, we mention here the curious fact that it is an outstanding open question of set theory whether one can find a model of ZFC that does not contain any higher Suslin line. Consequently there is a chance that theorem 7 yields us a ZFC example of a first countable GO space with no point-countable π -base.

4. EXAMPLES FROM SUBFAMILIES OF $\mathcal{P}(\omega)$

In this section we are going to introduce a (quite simple but apparently new) way of constructing first countable, 0-dimensional Hausdorff topologies on subfamilies of $\mathcal{P}(\omega)$, the power set of ω . Then we shall use some of the spaces obtained in this manner to present examples that demonstrate the necessity of the use of CH in Tkačenko's results mentioned in the introduction.

We start with fixing some notation and terminology. We shall use $\mathcal{U}$ to denote the family of all co-finite subsets of ω . For a given family

$\mathcal{I} \subset \mathcal{P}(\omega)$ and for $I \in \mathcal{I}$ and $U \in \mathcal{U}$ we put

$$[I, U]_{\mathcal{I}} = \{J \in \mathcal{I} : I \subset J \subset U\}.$$

If $\mathcal{I} = \mathcal{P}(\omega)$ then we shall omit the subscript.

Finally, we say that the family $\mathcal{I} \subset \mathcal{P}(\omega)$ is *stable* if $I \in \mathcal{I}$ and $I =^* J$ for $J \subset \omega$ imply $J \in \mathcal{I}$ as well. (Of course, here $I =^* J$ means that I and J are equal mod finite, i. e. their symmetric difference $I \Delta J$ is finite.)

Definition 8. Let us fix a family $\mathcal{I} \subset \mathcal{P}(\omega)$. We shall denote by $\tau_{\mathcal{I}}$ the topology on $\mathcal{I}$ generated by all sets of the form $[I, U]_{\mathcal{I}}$, where $I \in \mathcal{I}$ and $U \in \mathcal{U}$, and by $X_{\mathcal{I}}$ the space $\langle \mathcal{I}, \tau_{\mathcal{I}} \rangle$.

Of course, $X_{\mathcal{I}}$ is identical with the appropriate subspace of the maximal such space $X_{\mathcal{P}(\omega)}$. A few basic (pleasant) properties of the spaces $X_{\mathcal{I}}$ are given by the following proposition.

Proposition 9. *The spaces $X_{\mathcal{I}}$ are first countable, 0-dimensional and Hausdorff.*

Proof. Clearly, it suffices to show this for $\mathcal{I} = \mathcal{P}(\omega)$ because all three properties are inherited by subspaces.

Now, observe first that if $J \in [I, U] \cap [I', U']$ then

$$J \in [J, U \cap U'] \subset [I, U] \cap [I', U'],$$

hence the “intervals” $[I, U]$ form an open basis of $\tau_{\mathcal{P}(\omega)}$, moreover $\{[I, U] : I \subset U \in \mathcal{U}\}$ forms a countable neighbourhood base of the point I of $X_{\mathcal{I}}$.

Next, if $J \notin [I, U]$ then either $J \setminus U \neq \emptyset$ and then $[J, \omega] \cap [I, U] = \emptyset$, or $J \subset U$ and $I \setminus J \neq \emptyset$. But in the latter case we may pick an $n \in I \setminus J$ and then we have $J \subset U \setminus \{n\}$, moreover $[J, U \setminus \{n\}] \cap [I, U] = \emptyset$ because $n \in I$. This means that all basic open sets $[I, U]$ are also closed, that is $X_{\mathcal{P}(\omega)}$ is indeed 0-dimensional.

Finally, for every $I \in \mathcal{P}(\omega)$ we clearly have

$$\bigcap \{[I, U] : I \subset U \in \mathcal{U}\} = \{I\},$$

implying that $X_{\mathcal{P}(\omega)}$ is also Hausdorff. $\square$

For any family $\mathcal{I} \subset \mathcal{P}(\omega)$ we shall denote by $\text{cof}(\mathcal{I})$ the cofinality of the partial order $\langle \mathcal{I}, \subset \rangle$. Also, we say that a cardinal number κ is a *set caliber* of $\mathcal{I}$ if for every subfamily $\mathcal{J} \in [\mathcal{I}]^{\kappa}$ there are $\mathcal{K} \in [\mathcal{J}]^{\kappa}$ and $I \in \mathcal{I}$ such that $\cup \mathcal{K} \subset I$ or, less formally, among any κ -many members of $\mathcal{I}$ there are κ -many that have an upper bound in $\mathcal{I}$. We now connect these concepts concerning $\mathcal{I}$ with properties of the associated space $X_{\mathcal{I}}$.

Proposition 10. *For any subfamily $\mathcal{I} \subset \mathcal{P}(\omega)$ we have*

- (i) $d(X_{\mathcal{I}}) = \text{cof}(\mathcal{I}) \cdot \omega$;
- (ii) *if $\mathcal{I}$ is stable and κ is a cardinal with $\text{cf}(\kappa) > \omega$ then κ is a caliber of the space $X_{\mathcal{I}}$ if and only if κ is a set caliber of the family $\mathcal{I}$.*

Proof. The proof of (i) and the left-to-right direction of (ii) follows immediately from the fact that $\mathcal{K} \subset \mathcal{I}$ has an upper bound in $\mathcal{I}$ iff $\bigcap \{[I, \omega]_{\mathcal{I}} : I \in \mathcal{K}\} \neq \emptyset$. To see the other direction, assume that κ is a set caliber of the family $\mathcal{I}$ and consider a family $\mathcal{B}$ of κ -many basic open sets. Since $\text{cf}(\kappa) > \omega$ we may assume that $\mathcal{B} = \{[I, U) : I \in \mathcal{J}\}$ for $\mathcal{J} \in [\mathcal{I}]^{\kappa}$ and a fixed $U \in \mathcal{U}$. By our assumption there is a $K \in [\mathcal{J}]^{\kappa}$ which has an upper bound $K \in \mathcal{I}$. Clearly, then $K \cap U \in \mathcal{I}$ as $\mathcal{I}$ is stable and consequently

$$K \cap U \in \bigcap \{[I, U) : I \in \mathcal{K}\}.$$

□

After these preparatory propositions we can now present a result that will allow us to obtain nice examples of first countable spaces without point-countable π -bases.

Theorem 11. *Assume that $\mathcal{I} \subset \mathcal{P}(\omega)$ is stable, $\text{cof}(\mathcal{I}) > \omega$, and ω_1 is a set caliber of $\mathcal{I}$. Then $\pi \text{sw}(X_{\mathcal{I}}) > \omega$.*

Proof. Since $X_{\mathcal{I}}$ is first countable, and by (i) of proposition 10, we have

$$\pi(X_{\mathcal{I}}) = d(X_{\mathcal{I}}) = \text{cof}(\mathcal{I}) > \omega.$$

But, in view of part (ii) of proposition 10, ω_1 is a caliber of $X_{\mathcal{I}}$, consequently no π -base of $X_{\mathcal{I}}$ can be point-countable. □

Corollary 12. *Assume that there is a mod finite strictly increasing ω_2 -sequence in $\mathcal{P}(\omega)$. Then there is a first countable, 0-dimensional and Hausdorff space of cardinality ω_2 and having ω_1 as a caliber. In particular, MA_{ω_1} implies the existence of such a space.*

Proof. Let $\{A_{\alpha} : \alpha < \omega_2\} \subset \mathcal{P}(\omega)$ be a mod finite strictly increasing ω_2 -sequence, i. e. we have $|A_{\alpha} \setminus A_{\beta}| < \omega$ and $|A_{\beta} \setminus A_{\alpha}| = \omega$ whenever $\alpha < \beta < \omega_2$. It is obvious that the family

$$\mathcal{I} = \{I \subset \omega : \exists \alpha < \omega_2 \text{ with } I = {}^* A_{\alpha}\}$$

is stable and satisfies $|\mathcal{I}| = \text{cof}(\mathcal{I}) = \omega_2$. Next, we claim that $\mathcal{I}$ has ω_1 as a set caliber.

To see this, consider any family $\mathcal{J} = \{I_{\alpha} : \alpha \in a\} \subset \mathcal{I}$ where $a \in [\omega_2]^{\omega_1}$ and $I_{\alpha} = {}^* A_{\alpha}$ for all $\alpha \in a$ and pick $\beta < \omega_2$ such that $a \subset \beta$.

Then $|A_\alpha \setminus A_\beta| < \omega$ for all $\alpha \in a$, hence there is a fixed $s \in [\omega]^{<\omega}$ such that

$$b = \{\alpha \in a : A_\alpha \setminus A_\beta \subset s\}$$

is uncountable. But clearly, $s \cup A_\beta$ is then an upper bound of $\mathcal{K} = \{I_\alpha : \alpha \in b\}$ in $\mathcal{I}$. Consequently, by theorem 11, the space $X_{\mathcal{I}}$ is as required. $\square$

This result takes care of problems 4.6 and 4.7 from [8] by showing that it is consistent to have first countable Tychonov spaces with caliber ω_1 (and hence also CCC) without any point-countable π -base. With some further elaboration we may find examples that, in addition, are also hereditarily Lindelöf, and thus provide a solution to problem 4.3 as well.

Theorem 13. *Let $\{A_\alpha : \alpha < \omega_2\} \subset \mathcal{P}(\omega)$ be a mod finite strictly increasing ω_2 -sequence with the additional property that for every uncountable index set $a \in [\omega_2]^{\omega_1}$ there is a pair $\{\alpha, \beta\} \in [a]^2$ such that $A_\alpha \subset A_\beta$, (i. e. A_α is really a subset of A_β , not just mod finite). Then, for $\mathcal{I}$ defined as in corollary 12, the space $X_{\mathcal{I}}$ is also hereditarily Lindelöf.*

Proof. Assume, on the contrary, that $X_{\mathcal{I}}$ has an uncountable right-separated subspace. Without loss of generality this may be taken of the form $\{I_\alpha : \alpha \in a\}$, right separated in the natural well-ordering of its indices, where $a \in [\omega_2]^{\omega_1}$ and $I_\alpha =^* A_\alpha$ for all $\alpha \in a$. Moreover, we may assume that we have a fixed $U \in \mathcal{U}$ such that $[I_\alpha, U)_{\mathcal{I}}$ is a right separating neighbourhood of I_α for any $\alpha \in a$.

Clearly, there is a fixed finite set $s \in [\omega]^{<\omega}$ such that

$$b = \{\alpha \in a : I_\alpha \Delta A_\alpha = s\}$$

is uncountable. By our assumption, there is a pair $\{\alpha, \beta\} \in [b]^2$ (with $\alpha < \beta$) for which $A_\alpha \subset A_\beta$ and hence $I_\alpha \subset I_\beta$. This, however, would imply $I_\beta \in [I_\alpha, U)_{\mathcal{I}}$, contradicting that $[I_\alpha, U)_{\mathcal{I}}$ is a right separating neighbourhood of I_α . $\square$

Note that a space as in theorem 13 is a first countable L-space, hence unlike the spaces in corollary 12, it does not exist under MA_{ω_1} , see [7]. Luckily, there is a “natural” forcing construction that produces a mod finite strictly increasing ω_2 -sequence in $\mathcal{P}(\omega)$ with the additional property required in theorem 13.

Theorem 14. *There is a CCC forcing that, to any ground model, adds a mod finite strictly increasing sequence $\{A_\alpha : \alpha < \omega_2\} \subset \mathcal{P}(\omega)$ in any uncountable subsequence of which there are two members with proper inclusion.*

Proof. Let $\mathbb{P}$ consist of those finite functions $p \in Fn(\omega_2 \times \omega, 2)$ for which $\text{dom}(p) = a \times n$ with $a \in [\omega_2]^{<\omega}$ and $n < \omega$. We define $p' \leq p$ (i. e. p' extends p) as follows: $p' \supset p$, moreover $p'(\alpha, i) = 1$ implies $p'(\beta, i) = 1$ whenever $\alpha, \beta \in a$ with $\alpha < \beta$ and $i \in n' \setminus n$ (of course, here $\text{dom}(p) = a \times n$ and $\text{dom}(p') = a' \times n'$). It is straight-forward to show that $\langle \mathbb{P}, \leq \rangle$ is a CCC notion of forcing.

Let $G \subset \mathbb{P}$ be generic, then it follows from standard density arguments that $g = \cup G$ maps $\omega_2 \times \omega$ into 2 and if we set

$$A_\alpha = \{i < \omega : g(\alpha, i) = 1\}$$

then $\{A_\alpha : \alpha < \omega_2\}$ is mod finite strictly increasing.

To finish the proof, let us assume that $p \in \mathbb{P}$ forces that $\dot{h}$ is an order preserving injection of ω_1 into ω_2 . It will clearly suffice to show that p then has an extension q which forces $A_{\dot{h}(\xi)} \subset A_{\dot{h}(\eta)}$ for some $\xi < \eta < \omega_1$.

To see this, let us choose first for each $\xi < \omega_1$ a condition $p_\xi \leq p$ and an ordinal $\alpha_\xi < \omega_2$ such that $p_\xi \Vdash \dot{h}(\xi) = \alpha_\xi$. We may assume without any loss of generality that for some $n < \omega$ we have $\text{dom}(p_\xi) = a_\xi \times n$ and $\alpha_\xi \in a_\xi$ for all ξ . Using standard Δ -system and counting arguments, it is easy to find then $\xi < \eta < \omega_1$ such that p_ξ and p_η are compatible as functions and for any $i < n$ we have $p_\xi(\alpha_\xi, i) = p_\eta(\alpha_\eta, i)$. But then we clearly have $q = p_\xi \cup p_\eta \in \mathbb{P}$ and $q \leq p$, moreover it is obvious that q forces $A_{\alpha_\xi} \subset A_{\alpha_\eta}$ and hence $A_{\dot{h}(\xi)} \subset A_{\dot{h}(\eta)}$ as well. $\square$

From theorems 13 and 14 we immediately obtain a joint solution to problems 4.3 and 4.7 (and hence 4.6) of Tkačuk from [8].

Corollary 15. *It is consistent to have a first countable, hereditarily Lindelöf 0-dimensional T_2 space with caliber ω_1 having no point-countable π -base.*

Let us recall here that the failure of CH is not sufficient to produce a mod finite strictly increasing ω_2 -sequence in $\mathcal{P}(\omega)$, the basic ingredient of our examples in this section. In fact, Kunen had proved (see e. g. [6]) that if one adds ω_2 Cohen reals to a model of CH then no such sequence exists in the extension. Actually, we can show, as a strengthening of this, that in the same model if a subfamily $\mathcal{I}$ of $\mathcal{P}(\omega)$ has ω_1 as a set caliber then $\text{cof}(\mathcal{I}) \leq \omega$. So, we may not hope to use the methods of this section to find examples just assuming the negation of CH. So the following problem can be raised.

Problem 16. *Does $2^\omega > \omega_1$ imply the existence of a first countable Lindelöf and/or CCC Tychonov space having no point-countable π -base?*

REFERENCES

- [1] H. R. Bennett and T. G. McLaughlin, *A selective survey of axiom-sensitive results in general topology*, Texas Tech University Mathematics Series, No. 12. Lubbock, Tex., 1976. iv+114 pp.
- [2] A. Hajnal, *Proof of a conjecture of S. Ruziewicz*, Fund. Math. 50 (1961/1962), pp. 123–128.
- [3] A. Hajnal and I. Juhász, *On hereditarily α -Lindelf and hereditarily α -separable spaces*, Ann. Univ. Sci. Budapest, Sect. Math. 11 (1968), pp. 115–124.
- [4] I. Juhász, *Cardinal functions – 10 years later*, Math. Center Tract no. 123, Amsterdam, 1980
- [5] I. Juhász, *Cardinal functions II*, in: Handbook of Set Theoretic Topology, Eds: K. Kunen & J. E. Vaughan, Amsterdam, 1984, pp. 63–109.
- [6] I. Juhász, L. Soukup, and Z. Szentmiklóssy, *Combinatorial principles from adding Cohen reals*, Logic Colloquium '95 (Haifa), Lecture Notes in Logic 11, Springer, Berlin, 1998, pp. 79–103.
- [7] Z. Szentmiklóssy, *S-spaces and L-spaces under Martin's axiom*, Topology, Vol. II (Proc. Fourth Colloq., Budapest, 1978), Colloq. Math. Soc. Jnos Bolyai, 23, North-Holland, Amsterdam-New York, 1980, pp. 1139–1145.
- [8] V. V. Tkačuk, *Point-countable π -bases in first countable and similar spaces*, Fund. Math. 186 (2005), pp. 55–69.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS

E-mail address: juhasz@renyi.hu

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS

E-mail address: soukup@renyi.hu

EÖTVÖS LORÁNT UNIVERSITY, DEPARTMENT OF ANALYSIS, 1117 BUDAPEST,
PÁZMÁNY PÉTER SÉTÁNY 1/A, HUNGARY

E-mail address: zoli@renyi.hu